

## Effects of elevated atmospheric CO<sub>2</sub> on water relations of soya bean

Stephen A. Prior<sup>1</sup>, Hugo H. Rogers<sup>1</sup>, Nasser Sionit<sup>2</sup>, and Robert P. Patterson<sup>3</sup>

<sup>1</sup>USDA-ARS National Soil Dynamics Laboratory, Auburn, AL 36831-0792 (U.S.A.)

<sup>2</sup>Systems Ecology Research Group, College of Sciences, San Diego State University, San Diego, CA 92182-0057 (U.S.A.)

<sup>3</sup>Department of Crop Science, College of Agriculture and Life Sciences, North Carolina State University, Raleigh, NC 27695-7620 (U.S.A.)

(Accepted for publication 8 August 1990)

### ABSTRACT

Prior, S.A., Rogers, H.H., Sionit, N. and Patterson, R.P., 1991. Effects of elevated atmospheric CO<sub>2</sub> on water relations of soya bean. *Agric. Ecosystems Environ.*, 35: 13–25.

Soya bean (*Glycine max* (L.) Merr. 'Bragg') plants were grown in large containers in open-top field chambers under five atmospheric CO<sub>2</sub> concentrations (349–946  $\mu\text{l l}^{-1}$ ) and two water regimes. Rate of soil water depletion for the high CO<sub>2</sub> treatments started to decrease under well-watered conditions during anthesis and by early pod formation under water-stressed conditions. During reproductive growth, normal and stressed plants at 349  $\mu\text{l l}^{-1}$  (ambient level) received irrigation water 29 and 12 times, respectively, compared with 21 and 9 times, respectively, at 946  $\mu\text{l l}^{-1}$  CO<sub>2</sub>. At both anthesis and pod fill, plants grown under CO<sub>2</sub> enrichment exhibited greater leaf area. Nevertheless, water use per plant either remained constant (stressed plants at anthesis) or else declined (well-watered plants at anthesis; both moisture levels during pod fill) in response to CO<sub>2</sub> enrichment. At pod fill, leaves of CO<sub>2</sub>-enriched plants generally displayed a higher stomatal resistance, except near the end of the sampling period when a sudden increase in resistance was observed under low CO<sub>2</sub> owing to low soil water availability. Midday xylem potential for well-watered plants was greater than values for stressed plants and was unaffected by CO<sub>2</sub> treatment. Under low moisture conditions, elevated CO<sub>2</sub> had no effect on xylem potential at anthesis; however, during pod fill potential increased significantly with increasing CO<sub>2</sub> concentration, as elevated CO<sub>2</sub> decreased water use rates, lowering soil water stress. Alleviation of water stress during critical reproductive phases was strongly suggested.

### INTRODUCTION

The concentration of CO<sub>2</sub> in the atmosphere is known to be rising (Keeling et al., 1989), and if the current trend continues, a doubling of the present concentration ( $\sim 350 \mu\text{l l}^{-1}$ ) may occur during the next century (Gribbin, 1981). Although CO<sub>2</sub> is only a trace gas in the atmosphere, increases in its concentration could significantly affect global agriculture. Carbon dioxide is essential to plant growth as a source of carbon necessary for photosynthesis

and influences stomatal opening, hence affecting water vapor and gas exchanges. For a great many plant species, CO<sub>2</sub> enrichment is known to increase dry matter production and frequently increases net photosynthesis (Rogers et al., 1983a; Sionit et al., 1984; Strain and Cure, 1985; Allen et al., 1987; Warrick, 1988; Allen, 1990). Increased CO<sub>2</sub> levels induce stomatal closure, thereby decreasing transpiration rate which enhances water use efficiency (Carlson and Bazzaz, 1980; Kimball and Idso, 1983; Dahlman et al., 1985). On the other hand, the availability of soil water to crops is considered to be the major limitation to crop production in the U.S. (Boyer, 1982). Generally speaking, plant growth is reduced under soil water deficits owing to decreases in water potential, stomatal aperture, and photosynthesis (Boyer, 1970).

Consequently, research investigations have sought to evaluate the interactions of elevated CO<sub>2</sub> and limited soil water. Growth chamber studies with wheat (*Triticum aestivum* L.) have demonstrated that owing to the influence of CO<sub>2</sub> on water use efficiency (Gifford, 1979) and on water stress adaptation through osmoregulation (Sionit et al., 1981), enhancement of dry matter production under water stress conditions still occurs under CO<sub>2</sub> enrichment. Soya beans grown under CO<sub>2</sub> enrichment and subjected to water stress during early reproductive growth avoided the onset of severe water stress by exhibiting decreased water use (Rogers et al., 1984) and increased water use efficiency (Huber et al., 1984), thus enabling them to maintain higher leaf water potentials compared with those under low CO<sub>2</sub>. The results of these studies and the paucity of data on closely monitored long-term field studies on CO<sub>2</sub> and water stress interaction conducted where soil water was restricted throughout the soya bean growth cycle warrants further investigation. The objective of this study was to help fill that knowledge gap.

In the present work, soya bean plants were grown in large containers in open-top field chambers under various CO<sub>2</sub> levels and subjected to repeated drying cycles for the entire growing season. More specifically, the objectives of this investigation were : (a) to study the influence of these treatments on leaf water potential, stomatal resistance, and transpiration; (b) to document the influence of CO<sub>2</sub> concentration on water use of well-watered and water-stressed plants at two periods in reproductive growth; (c) to determine the relationship of leaf area and rate of water use at these same periods.

#### MATERIALS AND METHODS

Soya bean plants were grown from seeds in containers placed in open-top field chambers (Rogers et al., 1983b) at different atmospheric CO<sub>2</sub> concentrations and in open plots (i.e., no chambers) under ambient atmospheric conditions. Each chamber was constructed of a cylindrical aluminum frame (3 m in diameter by 2.4 m in height) covered with PVC film panels (Roll-A-

Glass\*) with a 45° frustum attached at the top; panels were held to the frame in farthingale fashion. Air with a predetermined amount of CO<sub>2</sub> was supplied continuously at 0.94 m<sup>3</sup> s<sup>-1</sup> through the perforated inner wall of the double-walled bottom half of each chamber. There were two replicates of six CO<sub>2</sub> treatments. Seasonal daytime CO<sub>2</sub> means ± 1 SD were: 348 ± 19 (ambient plots without chambers), 349 ± 19 (ambient chamber), 421 ± 22, 496 ± 24, 645 ± 33, and 946 ± 49 µl l<sup>-1</sup> (enriched chambers). A microcomputer was used to control the automatic delivery of air samples to a nondispersive infrared CO<sub>2</sub> analyzer, to log data, and to assist in regulating the injection of CO<sub>2</sub>.

Seeds were inoculated with commercial *Rhizobium* and planted on Julian Day 172 in large plastic containers (16.5 l) containing 13 l of a 2:1:1 mixture (by volume) of sandy clay loam soil: sand: \*Metro-Mix 220 (W.R. Grace Co.). Each study plot (chamber or open plot) contained 20 plants grown singly in containers which were randomly divided into two groups of 10 and subjected to two watering regimes. Differential water treatments were monitored with tensiometers, and rain was excluded by the erection of a removable transparent plastic covering during periods of rainfall. Containers were placed on racks to ensure proper drainage and exclude any groundwater. Plants were rewatered when tensiometers reached -0.005 to -0.015 MPa and -0.075 to -0.085 MPa for well-watered controls (WW; N=4) and water-stressed (WS; N=6) plants, respectively. Monitoring of drying cycles started when plants were at the third-node stage (V-3; Fehr and Caviness, 1977). Tensiometer readings (taken three times each day) and intervals between watering were recorded for the entire season. All tensiometers were flushed to eliminate air bubbles and refilled with degassed water each week. One liter of an N-free nutrient solution (Israel, 1981) was added to every container once a week.

The experimental design was a split plot with a randomized complete block arrangement of the main-plot factor (test atmospheres of CO<sub>2</sub>) for which there were two blocks. The treatments of the second factor (water treatments: WW and WS) were randomly assigned to subplots (containers) within each main plot.

Leaf transpiration, stomatal resistance, xylem pressure potential ( $\Psi_x$ ) and whole canopy transpiration (water use) measurements were made at two periods in reproductive growth: anthesis (RI) and pod fill (R5). All containers were watered to drip-point and WS plants were then allowed to go through a drying cycle while WW plants received irrigation water as needed. Corresponding drying cycles for RI and R5 occurred 55–61 and 94–102 days after planting (Fig. 1).

Leaf transpiration and stomatal resistance were determined with a LI-COR

\*Trade names and products are mentioned solely for information. No endorsement by the U.S. Department of Agriculture is implied.




Fig. 1. Changes in soil matric potential (A,B) during reproductive growth (55–94 days after planting) for well-watered and water-stressed soya bean grown at 349 (▲), 645 (■) and 946 (◆)  $\mu\text{l l}^{-1}$ . For WW,  $N=4$ ; for WS,  $N=6$ .

1600 Steady State Diffusive Resistance Porometer. At R5, daily measures were made on eight plants (four WW and four WS) per plot for both blocks. All measurements were obtained from the abaxial surface of fully expanded, sun-exposed trifoliates at the canopy top between 13:00 and 14:00 h.

A pressure chamber apparatus (PMS Instrument Co.) was used to measure xylem potential ( $\Psi_x$ ). At R1, readings were made at the end of the drying cycle. However, by R5, WS plants at the lower  $\text{CO}_2$  levels (348–496  $\mu\text{l l}^{-1}$ ) were depleting soil water faster than the high  $\text{CO}_2$  level plants; therefore, in order to make direct comparisons of plant water status at the various  $\text{CO}_2$  values, measurements of  $\Psi_x$  were made for all treatments prior to rewatering those with lower  $\text{CO}_2$ . Values of  $\Psi_x$  were taken from six plants (three WW and three WS) in each plot.

Water use was determined gravimetrically during the two sampling periods with a digital weighing scale. A total of six containers (three WW and three WS) from each plot were weighed every 24 h during the sampling periods. Leaf areas were obtained photometrically on these same plants. Thus, water use could then be expressed on both a per plant and a per unit leaf area basis.

Statistical analyses of data were performed according to standard analysis

of variance (ANOVA) and regression techniques using Statistical Analysis Systems (1985). All tests of statistical significance were conducted at  $P < 0.05$ .

## RESULTS AND DISCUSSION

Observations from three drying cycles during vegetative growth (data not shown) and the first two drying cycles in early reproductive phase (Fig. 1b) indicated that WS soya beans grown under the various CO<sub>2</sub> levels extracted soil water at similar rates. In addition, it was observed that vegetative drying cycles (data not shown) were approximately twice as long as those occurring in the early part of the reproductive period (Fig. 1b). Thus, there was a clear indication that soil water depletion had accelerated at the onset of reproduction. Similar observations have been made by other researchers studying soya bean under ambient levels of CO<sub>2</sub> (Teare et al., 1973; Kanemasu et al., 1976).

During reproductive growth, the positive effects of CO<sub>2</sub> enrichment on soil water conservations were observed earlier under the WW regimes than under the WS condition (Fig. 1a,b; see also Fig. 3a). Although high-CO<sub>2</sub> grown plants were larger, water extraction from the soil proceeded at a much lower rate. For example, under WS conditions on Day 70, the 349  $\mu\text{l l}^{-1}$  treatment reached the end of a drying cycle before the 645 or the 946  $\mu\text{l l}^{-1}$  treatment. High CO<sub>2</sub> plants encountered the same low water a few days later, but as the watering cycle was longer, these plants experienced fewer stressful days than low CO<sub>2</sub> plants over the season. Subsequent drying cycles were out of phase until re-watering on Day 94 (Fig. 2). Within this period, all plants were at the R5 stage. The initial drying cycle for both water treatments shown in Fig. 2b clearly demonstrates that plants grown under 645 and 946  $\mu\text{l l}^{-1}$  CO<sub>2</sub> had lower rates of soil water depletion than plants grown at 349  $\mu\text{l l}^{-1}$  CO<sub>2</sub>.

Overall, during reproductive growth, WS plants at 349  $\mu\text{l l}^{-1}$  CO<sub>2</sub> went through 12 drying cycles compared with 10 and 9 at 645 and 946  $\mu\text{l l}^{-1}$  CO<sub>2</sub>, respectively. Under WW conditions, plants grown at 349  $\mu\text{l l}^{-1}$  CO<sub>2</sub> received irrigation water 29 times compared with 24 and 21 times at 645 and 946  $\mu\text{l l}^{-1}$  CO<sub>2</sub>, respectively. Based on these observed values, we can infer that CO<sub>2</sub> enrichment has a strong tendency to alleviate plant water stress when critical periods of water demand occur during reproduction. Previous work has demonstrated that soya bean is more sensitive to water stress during pod and seed development than during vegetative growth (Sionit and Kramer, 1977; Korte et al., 1983). In order to examine this phenomenon at different levels of CO<sub>2</sub>, we examined in detail two periods during reproductive growth (R1 and R5).

Total leaf area and rates of water use per plant and per unit leaf area as affected by water regime and CO<sub>2</sub> concentration during two periods within the reproductive phase are shown in Figs. 3 and 4. At R1, water use per plant declined linearly with increasing CO<sub>2</sub> under WW conditions (Fig. 3a). Based on regression values from the 349 and 946  $\mu\text{l l}^{-1}$  treatments, we observed an




Fig. 2. Changes in soil matric potential (A,B) during reproductive growth (> 94 days after planting) for well-watered and water-stressed soya bean grown at 349 (▲), 645 (■) and 946 (◆)  $\mu\text{l l}^{-1}$ . For WW,  $N=4$ ; for WS,  $N=6$ .

approximate 15% reduction in plant water use. Decreases in water use per plant were documented even though plants grown under  $\text{CO}_2$  enrichment exhibited significant increases in leaf area (Fig. 3c). Leaf area was 27% greater at 946 than at 349  $\mu\text{l l}^{-1}$ . Thus, a more striking decrease in canopy transpiration was observed when water use was expressed on a unit leaf area basis (Fig. 3b). Rate of water use per unit leaf area was reduced by 47% at the 946  $\mu\text{l l}^{-1}$  treatment level compared with the 349  $\mu\text{l l}^{-1}$  control.

As expected, water stress significantly reduced rates of water use and leaf area at all levels of  $\text{CO}_2$ , relative to WW controls. More importantly, the rate of water use per plant was unaffected by  $\text{CO}_2$  treatment (Fig. 3a). In fact, plant water use remained relatively constant across the various  $\text{CO}_2$  treatment levels. These data verified observations from corresponding WS drying cycles which demonstrated that additional  $\text{CO}_2$  had no effect on reducing rates of soil water depletion (Days 55–61; Fig. 1b). On the other hand, leaf area increased linearly in response to elevated  $\text{CO}_2$  (Fig. 3c). Therefore, water use expressed on a unit leaf area basis exhibited a significant decrease (Fig. 3b). Water use per unit leaf area in plants at 946  $\mu\text{l l}^{-1}$   $\text{CO}_2$  was 27% less than in those of the 349  $\mu\text{l l}^{-1}$  control.

At R5, water use per plant decreased linearly for both water regimes in re-




Fig. 3. Water use per plant (A) and per unit leaf area (B) and leaf area per plant (C) for well-watered ( $\Delta$ ,  $\blacktriangle$ ) and water-stressed ( $\circ$ ,  $\bullet$ ) soya bean grown in open plots (open symbols) and five CO<sub>2</sub> treatments within chambers (closed symbols) at anthesis. Second term in the best fit linear regression equation represents significant chamber effect. Response to CO<sub>2</sub> was non-significant for WS in A.  $N=6$ .

sponse to increased CO<sub>2</sub> (Fig. 4a). These data reinforced findings from corresponding drying cycles which indicated that soil water extraction proceeded at a much lower rate under CO<sub>2</sub> enrichment (Days 94–102; Fig. 2b). We found no significant stress by CO<sub>2</sub> interaction, suggesting that plants in both water treatments had similar responses to additional CO<sub>2</sub>. Each water stress treatment reduced water use per plant by approximately the same proportion at each CO<sub>2</sub> level. Based on regression values from the 349 and 946  $\mu\text{L l}^{-1}$  treatment levels, water use per plant was reduced by about 32% at the high CO<sub>2</sub> level in both water treatments.

In terms of leaf area, we observed a highly significant stress by CO<sub>2</sub> interaction (Fig. 4c). Chronic water stress reduced leaf area. However, the adverse effects were most noticeable at the lower two treatment levels (349 and




Fig. 4. Water use per plant (A) and per unit leaf area (B) and leaf area per plant (C) for well-watered ( $\Delta$ ,  $\blacktriangle$ ) and water-stressed ( $\circ$ ,  $\bullet$ ) soya bean grown in open plots (open symbols) and five CO<sub>2</sub> treatments within chambers (closed symbols) at pod fill. Best fit linear and quadratic regression lines are shown. Chamber effect (second equation term) was significant in B and C. Response to CO<sub>2</sub> model was non-significant for WW in C.  $N=6$ .

421  $\mu\text{l l}^{-1}$ ). WW plants showed a slight but not statistically significant increase in leaf area as CO<sub>2</sub> was increased. In contrast, the leaf area of WS plants increased progressively with increasing CO<sub>2</sub> up to 645  $\mu\text{l l}^{-1}$ . Values of leaf area at both the 645 and 946  $\mu\text{l l}^{-1}$  steps were essentially the same. Leaf area was 32% higher in soya bean plants grown at 645  $\mu\text{l l}^{-1}$  CO<sub>2</sub> than in those grown at 349  $\mu\text{l l}^{-1}$ . However, when water use was expressed on a unit leaf area basis, CO<sub>2</sub> was the only significant source of variation (Fig. 4b). Rates of water use per unit leaf area declined in a nonlinear fashion with increasing CO<sub>2</sub>. Values at 645 and 946  $\mu\text{l l}^{-1}$  CO<sub>2</sub> were approximately 40% lower than at the 349  $\mu\text{l l}^{-1}$  control.

It is important to note that increased leaf area due to CO<sub>2</sub> under WS conditions (Fig. 3c) may have counteracted high stomatal resistance per unit leaf



area (data not shown). Nevertheless, whole canopy rate of water use (Fig. 3a) remained relatively unchanged with rising CO<sub>2</sub>. Similar results were obtained by Jones et al. (1984) working with soya bean at similar ages grown under optimal soil water conditions. These investigators found that diurnal bulk canopy resistance (stomatal plus boundary layer) was greater for high CO<sub>2</sub> soya bean canopies, but that this was offset by a larger transpiring leaf area and higher vapor pressure deficits. They further observed that absolute water use was approximately the same across various CO<sub>2</sub> concentrations. Likewise, in a growth chamber study, Gifford (1979) reported little change in the actual amount of water used by wheat grown under CO<sub>2</sub> enrichment and different water regimes. Conversely, in agreement with observations reported by Rogers et al. (1984), the increase in leaf area (Fig. 3c) under WW conditions was apparently not sufficient to offset the high stomatal resistance per unit area (data not shown), thus total canopy water use rate decreased (Fig. 3a) with increasing CO<sub>2</sub> concentration.

Figure 5 illustrates the effects of CO<sub>2</sub>, and water treatment on leaf stomatal resistance and transpiration over several days at R5. As reported elsewhere for other species (Carlson and Bazzaz, 1980; Patterson and Flint, 1982), throughout the sampling period, leaves of CO<sub>2</sub> enriched plants grown under WW conditions had higher stomatal resistance (Fig. 5a) and lower transpir-




Fig. 5. Leaf stomatal resistance (A,C) and transpiration (B,D) for well-watered (A,B) and water-stressed (C,D) soya bean grown in five CO<sub>2</sub> treatments within chambers (●, 349; ◆, 421; ●, 496; ▲, 645; ■, 946 μl l<sup>-1</sup>) at pod fill. Measurements were taken for 6 days during a drying cycle (Days 94–102). *N* = 8.

ation rates (Fig. 5b) relative to those WW plants in ambient  $\text{CO}_2$ . Under WS conditions, the stomatal response at 645 and 946  $\mu\text{l l}^{-1}$   $\text{CO}_2$  (Fig. 5c) resulted in transpiration rates (Fig. 5d) which were similar to those observed under WW conditions (Fig. 5b). Even during early portions of stress periods, lower levels of elevated  $\text{CO}_2$  (421 and 496  $\mu\text{l l}^{-1}$ ) induced partial stomatal closure which caused lower transpiration rates relative to the 349  $\mu\text{l l}^{-1}$  control. However, from Day 3 onward, soya beans grown at 349–496  $\mu\text{l l}^{-1}$   $\text{CO}_2$ , exhibited similar decreases in transpiration rate in response to decreasing water availability. Likewise, the sudden rise in stomatal resistance observed at 349–496  $\mu\text{l l}^{-1}$   $\text{CO}_2$  on Day 6 (Fig. 5d) reflects plant response (i.e., stomatal closure) to low soil water availability. Similar to observations reported by Rogers et al. (1984), transpiration at all  $\text{CO}_2$  levels fell to nearly the same low rate on this day despite the large differences in resistance.

Midday  $\Psi_x$  data from both reproductive periods are shown in Fig. 6. At R1,  $\text{CO}_2$  enrichment had little effect on reducing  $\Psi_x$  under water stress (Fig. 6a). The fact that similar midday stresses were seen was not surprising since plants at the various  $\text{CO}_2$  concentrations had extracted soil water at equivalent rates (Fig. 3a) and therefore reached approximately the same soil moisture level by the time of measurement (Day 60; Fig. 1b). For WW plants, there were no significant differences in  $\Psi_x$  due to  $\text{CO}_2$ . Comparable results have been reported for soya bean, sicklepod (*Cassia obtusifolia* L.), and showy crota-




Fig. 6. Leaf xylem potential for well-watered ( $\Delta$ ,  $\blacktriangle$ ) and water-stressed ( $\circ$ ,  $\bullet$ ) soya bean grown in open plots (open symbols) and five  $\text{CO}_2$  treatments within chambers (closed symbols) at anthesis (A: Day 60) and pod fill (B: Day 94). Best fit lines are shown. Chamber effect was not significant. Response to  $\text{CO}_2$  was non-significant in A and for WW in B.  $N=6$ .

laria (*Crotalaria spectabilis* Roth) measured 35 days after planting (Patterson and Flint, 1982). A significant stress effect was detected; however, the actual magnitudes of  $\Psi_x$  for WS plants were not much lower than WW plants, the differences in soil water notwithstanding. Evaporative demands at this sampling period were such that any difference in  $\Psi_x$  due to soil water or CO<sub>2</sub> level should have been detected. Therefore, the small observed differences in  $\Psi_x$  were probably due to plant size and soil water deficits being too low to cause a substantial difference in plant water deficit at this growth stage (Sionit and Kramer, 1976).

At R5 we observed a well-defined water stress by CO<sub>2</sub> interaction (Fig. 6b). There was little difference in  $\Psi_x$  due to CO<sub>2</sub> under WW conditions. In contrast, under WS conditions,  $\Psi_x$  increased significantly with rises in CO<sub>2</sub>. This response is confirmed when one recalls that WS plants grown under elevated CO<sub>2</sub> had lower rates of water use (Fig. 4a) and thus were under a lower level of stress (Fig. 2b). Plants grown at 349, 421 and 496  $\mu\text{l l}^{-1}$  CO<sub>2</sub> had  $\Psi_x$  values of  $-1.08$ ,  $-0.83$ , and  $-0.72$  MPa, respectively. Although Boyer (1970) and Ghorashy et al. (1971) reported that leaf water potentials below  $-1.1$  MPa were necessary to inhibit photosynthesis of young soya bean leaves, the values of  $\Psi_x$  attained by WS plants grown at these CO<sub>2</sub> levels were probably low enough to inhibit afternoon photosynthesis as wilting (as evidenced by severe flagging of leaves) was observed at these treatment levels. Wilted leaves recovered completely after rewatering. On the other hand, as the soil water levels were still relatively high (Fig. 2b) under WS conditions at the 645 and 946  $\mu\text{l l}^{-1}$  levels, their respective  $\Psi_x$  values were similar to their WW control values (Fig. 6b). Likewise, others have reported that the water potential of soya bean (Rogers et al., 1984) and wheat (Sionit et al., 1981) growing under low CO<sub>2</sub> concentrations declined more rapidly and reached a lower value at the end of the stress period relative to those grown under high CO<sub>2</sub>. In the present study, high CO<sub>2</sub> plants showed no signs of wilting at this time or when the plants reached a similar low soil water level a few days later (data not shown). Ambient grown plants showed extensive wilting at a time when other environmental factors are most conducive to rapid carbon fixation; thus, plants grown at high CO<sub>2</sub> presumably have an advantage over plants grown at a lower CO<sub>2</sub> level.

The results from this field experiment (conducted with soya bean plants grown in large containers) strongly indicates that the benefit of additional CO<sub>2</sub> on reducing the rate of water consumption occurs primarily at the beginning of reproductive growth (R1–R2) under WW conditions, but is delayed until the R3 stage under WS conditions. Detailed measurements from two stages in reproductive growth (R1 and R5) clearly showed that larger plants produced as a result of CO<sub>2</sub> enrichment generally had more leaf surface area. However, this was not sufficient to counteract the lower rate of transpiration per unit leaf area. Water use per plant either remained constant (WS plants

at R1) or declined (WW plants at R1; WW and WS plants at R5) in response to CO<sub>2</sub> enrichment. Our results suggest that a decreased rate of water use could, in itself, alleviate water deficit in soya bean at critical periods of water demand during reproductive growth. Reduced water use coupled with increases in dry matter production as also reported elsewhere for soya bean (Rogers et al., 1986) and other plants (Kimball, 1983) could lead to increases in water use efficiency which would have a substantial impact on agricultural productivity. However, further research with various economically important species grown in conventional row systems and under higher levels of soil water stress will be required in order to make accurate predictions concerning the potential effects of elevated atmospheric CO<sub>2</sub> on crop water use.

#### ACKNOWLEDGMENTS

This work was funded by the Department of Energy through Interagency No. DEA-01-81-ERG0001 to the USDA-ARS. We thank the ARS Air Quality Program (Raleigh, NC) for support facilities and for the help of their staff.

#### REFERENCES

- Allen, L.H., Jr., 1990. Plant responses to rising carbon dioxide and potential interactions with air pollutants. *J. Environ. Qual.*, 19: 15–34.
- Allen, L.H., Jr., Boote, K.J., Jones, J.W., Jones, P.H., Valle, R.R., Acock, B., Rogers, H.H. and Dahlman, R.C., 1987. Response of vegetation to rising carbon dioxide: photosynthesis, biomass, and seed yield of soybean. *Global Biogeochem. Cycles*, 1 (1): 1–14.
- Boyer, J.S., 1970. Differing sensitivity of photosynthesis to low leaf water potentials in corn and soybeans. *Plant Physiol.*, 46: 236–239.
- Boyer, J.S., 1982. Plant productivity and environment. *Science*, 218: 443–448.
- Carlson, R.W. and Bazzaz, F.A., 1980. The effects of elevated CO<sub>2</sub> concentration on growth, photosynthesis, transpiration, and water use efficiency of plants. In: J. Singh and A. Deepak (Editors), *Environmental and Climatic Impact of Coal Utilization*. Academic Press, New York, pp. 609–612.
- Dahlman, R.C., Strain, B.R. and Rogers, H.H., 1985. Research on the response of vegetation to elevated atmospheric carbon dioxide. *J. Environ. Qual.*, 14: 1–8.
- Fehr, W.R. and Caviness, C.E., 1977. Stages of soybean development. *Iowa Agric. Home Econ. Exp. Stn., Iowa Coop. Ext. Serv. Spec. Rep.* 80.
- Ghorashy, S.R., Pendleton, J.W., Peters, D.B., Boyer, J.S. and Beuerlein, J.E., 1971. Internal water stress and apparent photosynthesis with soybeans differing in pubescence. *Agron. J.*, 63: 674–676.
- Gifford, R.M., 1979. Growth and yield of CO<sub>2</sub>-enriched wheat under water limited conditions. *Aust. J. Plant Physiol.*, 6: 367–378.
- Gribbin, J., 1981. The politics of carbon dioxide. *New Sci.*, 90: 82–84.
- Huber, S.C., Rogers, H.H. and Mowry, F.L., 1984. Effects of water stress on photosynthesis and carbon partitioning in soybean [*Glycine max* (L.) Merr.] plants grown in the field at different CO<sub>2</sub> levels. *Plant Physiol.*, 76: 244–249.

- Israel, D.W., 1981. Cultivar and *rhizobium* strain effects on nitrogen fixation and remobilization by soybeans. *Agron. J.*, 73: 509–516.
- Jones, P., Allen, L.H., Jr., Jones, J.W., Boote, K.T. and Campbell, W.J., 1984. Soybean canopy growth, photosynthesis, and transpiration responses to whole-season carbon dioxide enrichment. *Agron. J.*, 76: 633–637.
- Kanemasu, E.T., Stone, L.R. and Powers, W.L., 1976. Evapotranspiration model tested for soybean and sorghum. *Agron. J.*, 68: 569–572.
- Keeling, C.D., Bacastow, R.B., Carter, A.F., Piper, S.C., Whorf, T.P., Heimann, M., Mook, W.G. and Roeloffzen, H., 1989. A three dimensional model of atmospheric CO<sub>2</sub> transport based on observed winds: observational data and preliminary analysis. Appendix A. In: Aspects of climate variability in the Pacific and the western Americas. *Geophys. Mono. Am. Geophys. Union*, Vol. 55, Nov.
- Kimball, B.A., 1983. Carbon dioxide and agricultural yield: An assemblage and analysis of 430 prior observations. *Agron. J.*, 75: 779–788.
- Kimball, B.A. and Idso, S.B., 1983. Increasing atmospheric CO<sub>2</sub>: Effects on crop yield, water use and climate. *Agric. Water Manage.*, 7: 55–72.
- Korte, L.L., Specht, J.E., Williams, J.H. and Sorensen, R.C., 1983. Irrigation of soybean genotypes during reproductive ontogeny. II. Yield component responses. *Crop Sci.*, 23: 528–533.
- Patterson, D.T. and Flint, E.P., 1982. Interacting effects of CO<sub>2</sub> and nutrient concentration. *Weed Sci.*, 30: 389–394.
- Rogers, H.H., Bingham, G.E., Cure, J.D., Smith, J.M. and Surano, K.A., 1983a. Responses of selected plant species to elevated CO<sub>2</sub> in the field. *J. Environ. Qual.*, 12: 569–574.
- Rogers, H.H., Heck, W.W. and Heagle, A.S., 1983b. A field technique for the study of plant responses to elevated carbon dioxide concentrations. *J. Air Pollut. Control Assoc.*, 33: 42–44.
- Rogers, H.H., Sionit, N., Cure, J.D., Smith, J.M. and Bingham, G.E., 1984. Influence of elevated carbon dioxide on water relations of soybeans. *Plant Physiol.*, 74: 233–238.
- Rogers, H.H., Cure, J.D. and Smith, J.M., 1986. Soybean growth and yield response to elevated carbon dioxide. *Agric. Ecosystems Environ.*, 16: 193–204.
- Sionit, N. and Kramer, P.J., 1976. Water potential and stomatal resistance of sunflower and soybean subjected to water stress during various growth stages. *Plant Physiol.*, 58: 537–540.
- Sionit, N. and Kramer, P.J., 1977. Effect of water stress during different stages of growth of soybean. *Agron. J.*, 69: 274–278.
- Sionit, N., Strain, B.R., Hellmers, H. and Kramer, P.J., 1981. Effects of atmospheric CO<sub>2</sub> concentration and water stress on water relations of wheat. *Bot. Gaz.*, 142: 191–196.
- Sionit, N., Rogers, H.H., Bingham, G.E. and Strain, B.R., 1984. Photosynthesis and stomatal conductance with CO<sub>2</sub>-enrichment of container and field-grown soybeans. *Agron. J.*, 76: 447–451.
- Statistical Analysis Systems, 1985. SAS User's Guide. SAS Institute, Cary, NC.
- Strain, B.R. and Cure, J.D. (Editors), 1985. Direct effect of increasing carbon dioxide on vegetation. Carbon dioxide research, state of the art. U.S. Department of Energy, Office of Energy Research, Washington, DC.
- Teare, I.D., Kanemasu, E.T., Powers, W.L. and Jacobs, H.S., 1973. Water-use efficiency and its relation to crop canopy area, stomatal regulation, and root distribution. *Agron. J.*, 65: 207–211.
- Warrick, R.A., 1988. Carbon dioxide, climatic change and agriculture. *Geogr. J.*, 154(2): 221–233.